

Minireview

Custom astrocyte-mediated vasomotor responses to neuronal energy demand

Jillian L LeMaistre and Christopher M Anderson

Address: Department of Pharmacology and Therapeutics, University of Manitoba, and Division of Neurodegenerative Disorders, St Boniface Hospital Research Centre, 351 Taché Avenue, Winnipeg, MB R2H 2A6, Canada.

Correspondence: Christopher M Anderson. Email: canderson@sbr.ca

Published: 16 February 2009

Genome Biology 2009, **10**:209 (doi:10.1186/gb-2009-10-2-209)

The electronic version of this article is the complete one and can be found online at <http://genomebiology.com/2009/10/2/209>

© 2009 BioMed Central Ltd

Abstract

Astrocytes mediate either constriction or dilation of local brain arterioles in response to synaptic activity. Recent work indicates that the directionality of this response may be dictated by ambient oxygen levels.

In the brain, astrocyte processes are arranged in coordinated non-overlapping spatial domains such that the vast majority of the surface area of cerebral arterioles and capillaries is contacted by astrocyte endfeet [1]. This makes astrocytes uniquely well positioned to send vasoactive signals to the blood-brain barrier. Several studies now present a consensus view that astrocytes respond to input from glutamate-producing (glutamatergic) neurons by increasing intracellular Ca^{2+} , phospholipase A2 (PLA2)-mediated arachidonic acid formation, and metabolism of arachidonic acid to produce either vasodilatory [2-6] or vasoconstrictor [5,7] metabolites. The conditions under which astrocytes elicit vasodilation or vasoconstriction have remained largely confusing, as either domination of one effect over the other [2,4,7,8] or both effects have been reported in various tissue slice models [5]. In addition, it has not been clear how astrocyte participation in functional or hypoxic hyperemia (an increase in blood flow) fits with the involvement of metabolic mediators such as adenosine, lactate, H^+ , K^+ and CO_2 , all of which have been implicated in this critical neurophysiological process [9,10]. A recent report from the laboratory of Brian MacVicar (Gordon *et al.* [11]) significantly clarifies both these issues. Here we discuss this work and give an overview of the multiple pathways by which astrocytes influence neuronal energy supply.

Do astrocytes cause vasodilation or vasoconstriction?

Zonta *et al.* [2] provided the first direct demonstration that astrocytes in brain slices sense increased levels of glutamate (due to neuronal activity) via metabotropic glutamate receptors (mGluRs) and respond with an increase in endfoot Ca^{2+} , activation of PLA2 and metabolism of arachidonic acid by COX-1 to produce vasodilatory prostaglandins (probably PGE_2) [2,12]. Subsequent studies confirmed a vasoactive role for astrocyte mGluR activation and Ca^{2+} elevation, but reported vasoconstriction rather than vasodilation [7]; constriction was attributed to PLA2-mediated arachidonic acid production by astrocytes, followed by diffusion of the arachidonic acid to smooth muscle and subsequent metabolism by cytochrome P450 4A (ω -hydroxylase) to 20-hydroxy-eicosatetraenoic acid (20-HETE). The clear objective left in the wake of these initial papers was to determine whether constriction or dilation was the 'physiological' response to astrocyte activation. Dilation required the presence of a nitric oxide synthase (NOS) inhibitor or other pre-constrictor [2,5-7], and was converted to constriction when NOS inhibitors were excluded [7], raising the possibility that dilations could only be produced by artificial enhancement of baseline vascular tone. On the other hand, constriction was intuitively less attractive as the natural response to increased neuronal work. More recent *in vivo* studies in

anesthetized mice provided strong support for dilation by showing that increased astrocyte Ca^{2+} levels produced only COX-1-dependent vasodilations and corresponding local increases in blood flow [3]. This was a crucial observation as it verified astrocyte-mediated control of cerebral microcirculation - specifically dilations - outside the brain slice model, which can be critiqued as non-physiological due to the lack of arterial pressurization. It was also shown recently that inhibition of 20-HETE synthesis failed to affect integrated somatosensory hyperemic vasodilation [13], providing further evidence against a physiological role for constrictions.

This dominant vasodilatory effect of astrocyte Ca^{2+} signaling *in vivo* called into question the practical significance of the astrocyte- and 20-HETE-mediated constrictions observed in brain slices. Some clarification came from studies showing that both vasodilation and vasoconstriction are possible, depending on the conditions employed [5,14]. One such study found that the polarity of astrocyte vasomotor influence is dependent on pre-existing smooth-muscle tone [14]. In isolated retinas, vasodilations dependent on astrocyte metabolism of arachidonic acid by cytochrome (CYP) P450 2C11 (epoxygenase) to epoxyeicosatrienoic acids (EETs) became less likely as NO levels increased and directly inhibited epoxygenase activity [5]. This identified NO as a pro-constriction factor despite its well-known actions as a direct cGMP-dependent vasodilator [5,15]. These results could explain discrepancies between earlier observations of vasodilations, in the presence of a NOS inhibitor [2], and vasoconstrictions, in conditions of no pre-constriction that probably included higher endogenous NO levels [7]. An opposite pro-dilatory effect of NO has also been observed owing to NO-mediated inhibition of CYP P450 4A and 20-HETE production [13] and it is not yet clear how these effects of NO at CYP P450s are regulated.

The report by Gordon *et al.* [11] significantly improves our understanding of the balance between vasoconstriction and vasodilation in brain slices. They discovered that although the mGluR-induced rise in astrocyte Ca^{2+} led to 20-HETE-dependent vasoconstriction in slices maintained in high O_2 (95%), the same treatment yielded PGE₂-dependent vasodilation in slightly hypoxic medium (20% O_2). The mechanism for this functional switch is twofold: first, mGluR activation increased the rate of glycolysis in astrocytes, leading to increased extracellular lactate levels sufficient to inhibit PGE₂ reuptake via the PGE₂/lactate exchanger (PGT). This resulted in a net increase in extracellular PGE₂ and a shift in the vasomotor balance from vasoconstriction towards vasodilation. Second, the authors postulate that the final tipping point from dilation to constriction is the action of adenosine at smooth muscle A_{2A} receptors, which overrides the 20-HETE-mediated constrictor effect. They showed that adenosine is produced in low O_2 , that exogenous adenosine can block 20-HETE-induced vasoconstriction at high (95%) O_2 , and that

mGluR-mediated vasoconstrictions can be converted to dilations by adding exogenous adenosine and PGE₂.

Overall, the findings of Gordon *et al.* [11] effectively establish that astrocytes mediate bidirectional control of local arteriolar diameter in a manner dictated by tissue metabolic status. In this light, one can envisage a 'see-saw'-like balance between vasodilation and vasoconstriction with weight for vasoconstriction provided by 20-HETE and intracellular astrocyte NO, and weight on the vasodilation end provided by adenosine, lactate and PGE₂ and/or EETs. It remains to be determined what fulcrum O_2 level must be reached before vasodilation is preferred. It is fairly certain that this level is in the normoxic O_2 range as *in vivo* functional hyperemia paradigms at normal PO_2 invariably produce dilation and increased cerebral blood flow [3], and the current findings by Gordon *et al.* [11] show that vasodilation dominated near the low end of physiological PO_2 in brain slices. Determination of where the switch point for dilation is may rest with characterizing how low PO_2 must fall before ATP production is compromised and adenosine accumulates. While Gordon *et al.* [11] demonstrated adenosine production in the low physiological range of PO_2 , it seems unlikely that slice PO_2 in the mid to high normoxic range (30-50 mmHg) would produce enough adenosine to maintain dilatory efficacy. At this point, one might predict that vasoconstriction would begin to prevail. Alternatively, it remains possible that local transient increases in extracellular adenosine resulting from breakdown of extracellular ATP during neuro- or gliotransmission could be sufficient to occupy A_{2A} receptors and drive vasodilation independent of a direct effect of PO_2 .

New ideas about 'old' vasodilators

The paper by Gordon *et al.* [11] provides valuable hints about how metabolic intermediates, including lactate, H^+ , K^+ and adenosine [9,16,17], might participate in a more regulated and coordinated hyperemic response than would be allowed simply by the diffusion of accumulated metabolites. The authors show that lactate and adenosine, in particular, participate in shifting the astrocyte vasomotor balance from constriction to dilation.

Lactate has direct vasodilatory effects *in vitro* [17] and augments increases in cerebral blood flow dependent on neuronal activity [10], but Gordon *et al.* [11] have identified a completely novel mechanism by which it can participate in vasodilation. They confirm previous observations [18] that mGluR activation drives astrocyte glycolysis and produces lactate in low O_2 , and also show that preventing the conversion of pyruvate to lactate eliminated mGluR-mediated vasodilation in brain slices, making the first specific link between astrocyte lactate and vasodilation. More importantly, they showed that the dilatory effects of lactate are

mediated indirectly by PGE₂, which is enhanced when extracellular lactate interferes with PGT-mediated exchange of intracellular lactate for extracellular PGE₂. These findings establish lactate as a novel operator of the astrocyte vasomotor switch and are consistent with the ability of lactate to augment neuronal-activity-induced increases in blood flow (modeled by Gordon *et al.* by using mGluR activation) without affecting resting blood flow [10].

It should be noted that while the authors provide evidence that mGluR-driven glycolysis in astrocytes is a source of lactate that can inhibit PGT and enhance extracellular PGE₂, the contribution of astrocyte lactate produced by other mechanisms may also be significant. For example, astrocyte lactate can be derived from intracellular Na⁺ accumulated during glutamate uptake [19]. A significant portion of the hyperemic response in olfactory bulb was recently shown to be dependent on glutamate transport, suggesting that this pathway may represent an important source of lactate [4]. Glycogen mobilization also drives astrocyte lactate production [20] but the contribution of this pathway is not known.

Adenosine acts as an inhibitory neuromodulator in the central nervous system and is implicated in regulating cerebral arterial tone in periods of increased neuronal activity [21] and in hyperemia precipitated by hypoxia [22] and hypoglycaemia [23]. There is also strong evidence for a direct dilatory effect of adenosine at vascular A2A receptors during hyperemia [24]. Gordon *et al.* [11] showed that an A2A receptor agonist blocked the mGluR-induced vasoconstriction normally observed at 95% O₂. They also converted constrictions to dilations in the same high O₂ conditions by combining exogenous adenosine with a PGT blocker, which mimicked the increased PGE₂ levels observed during lactate production at lower O₂ levels. These observations support previous reports that A2A receptors are vasodilatory in hyperemia and indicate that multiple vasomotor effects are additive; in other words, A2A-mediated dilation is capable of canceling 20-HETE-induced vasoconstriction.

The experiments of Gordon *et al.* [11] raise at least three intriguing questions on the role of adenosine in hyperemia. First, the authors nicely demonstrate that exogenous adenosine can compete with 20-HETE-mediated vasoconstriction in conditions of high O₂ but do not test whether endogenous adenosine is required to yield dilations at 20% O₂. It would be interesting to investigate whether mGluR-mediated vasodilation at 20% O₂ is sensitive to inhibition by blocking the effects of endogenous adenosine with an A2A receptor antagonist or exogenous adenosine deaminase. The current evidence leaves open the possibility that accumulation of PGE₂ by lactate-mediated inhibition of PGT is sufficient to overcome the effects of 20-HETE and produce dilation without endogenous adenosine. Second, while Gordon *et al.* [11] undoubtedly show that A2A receptor effects can add to

the vascular effects of 20-HETE and PGE₂, leading to dilation, it remains to be seen whether other metabolic vasodilators can contribute in the same additive way. It is not yet known, for example, whether H⁺, K⁺ or the direct vasodilatory effects of lactate itself can or do compete with the effects of 20-HETE. Lastly, Gordon *et al.* [11] confirm the widely held consensus that A2A adenosine receptors are likely to play an important role in hyperemia. Given the revelation that astrocyte A2B adenosine receptors regulate EET production and neurovascular coupling *in vivo* [25], it will be important to dissect the effects of endogenous adenosine to determine whether adenosine acts both at vascular A2A and astrocyte A2B receptors to influence the vasomotor balance toward dilation.

The contribution of Gordon *et al.* [11] together with that of Metea and Newman [5] show that previously identified vasoactive mediators such as NO, adenosine and lactate may converge on a central control point by influencing an astrocyte-controlled vasomotor balance. The result is a clearer view of the interplay among vasoactive effectors and a conceptually tantalizing model of coordinated spatial cerebral blood flow regulation by targeted vasodilations and vasoconstrictions mediated by Ca²⁺ signal propagation in distinct astrocyte networks.

Astrocytes coordinate a multimodal nutritive response to challenged neurons

The findings that astrocyte vasomotor polarity depends on oxygen-influenced changes in adenosine and activity-driven extracellular lactate concentrations are novel and exciting but do not exist in isolation. Rather, they are part of a multifaceted response by astrocytes to neuronal energy demand that also involves direct shuttling of tricarboxylic acid (TCA) cycle carbon sources from astrocytes to neurons (Figure 1). Perivascular astrocytes react in several ways to glutamatergic signals from neurons. Glutamate released by active neurons is rapidly removed from the synapse by high-affinity astrocyte glutamate transporters [26] and the internalized glutamate is converted to glutamine, which is released by astrocytes and claimed by neurons. This enables neurons to avoid a net loss of glutamate during neurotransmission and provides a potential TCA substrate for synthesis of γ -aminobutyric acid and production of ATP. Lactate is also shuttled from astrocytes to neurons for use as an oxidative fuel. Lactate increases have been reported in response to mGluR activation [18] and Na⁺-dependent glutamate transporter activity [19], and lactate can also be derived from mobilization of astrocyte glycogen in response to neuronal activity [20,27]. Intracellular lactate is released from astrocytes via the monocarboxylate transporter-1 (MCT1) where it can be taken up by neuronal MCT2 and converted to pyruvate for use in the TCA cycle. Importantly, astrocyte-neuron metabolic communication, and probably vasomotor communication, cannot be viewed simply as




Figure 1

Astrocyte influences on neuronal energy supply. Perivascular astrocytes respond to neuronal input (activity) by supplying neurons with substrates for oxidative phosphorylation (lactate, glutamine (Gln)) and glutamate (Glu) replenishment (glutamine), and by signaling changes in local blood flow at the vascular level. Active neurons produce synaptic glutamate that can be taken up by astrocyte glutamate transporters (EAAT) or activate mGluRs. (1) EAAT activation drives electrogenic Na⁺ influx, activates Na⁺/K⁺ ATPases and stimulates glycolytic lactate generation. (2) mGluR activation also leads to glycolysis and lactate production, and neuronal activity drives astrocyte glycogenolysis (3) and eventual lactate formation. Lactate from these three sources is released to the extracellular space via monocarboxylate transporter 1 (MCT1) where it can be taken up by neuronal MCT2 and converted to pyruvate (Pyr) for entry into the TCA cycle (4). Glutamate taken up by astrocyte EAATs can also be converted to glutamine by glutamine synthetase (5). Glutamine can be released and taken up by neuronal amino acid transporters for re-synthesis of glutamate and/or γ -aminobutyric acid via the TCA cycle. For astrocyte changes in blood flow, mGluR activation causes increased Ca²⁺ levels (6), leading to phospholipase A2 (PLA2) activation, arachidonic acid (AA) formation (7) and vasoconstriction following 20-HETE production by cytochrome P450 ω -hydroxylase (8) and continuous prostaglandin E₂ (PGE₂) generation by cyclooxygenase (COX) (9). Vasodilation can result in hypoxic conditions from lactate-mediated inhibition of PGE₂ clearance by prostaglandin transporters (PGT) following PGE₂ diffusion to the vascular smooth muscle (10). EAAT, excitatory amino acid transporter; Pyr, pyruvate.

signaling between discrete cell pairs or even among small groups of cells. New information affirms that lactate from cerebral arterioles is delivered over large distances along gap-junction-coupled astrocyte networks to areas of stimulated neurons but not to resting neurons [28]. Overall, astrocytes mount a multi-pronged neuronal aid effort centered largely on the multiple vasomotor and metabolic effects of lactate.

Studies over the past five years have revealed that astrocytes link neuronal energy supply and demand by triggering adaptive changes in the delivery of blood-borne glucose and O₂ to neurons. Gordon *et al.* [11] have taken the understanding of this to a new level by showing that astrocytes can act as switches to either increase or decrease blood flow to working neurons depending on regional metabolic status. Their study will serve as an important

launch point for future work aimed at identifying how astrocyte networks regulate the spatial control of brain blood flow both near to and distant from areas of neuronal activation.

Acknowledgements

We thank the Canadian Institute of Health Research and Manitoba Health Research Council for research support. CMA is supported by the Heart and Stroke Foundation of Canada. JLM is supported by a doctoral research award from the Canadian Institutes of Health Research.

References

- Nedergaard M, Ransom B, Goldman SA: **New roles for astrocytes: redefining the functional architecture of the brain.** *Trends Neurosci* 2003, **26**:523-530.
- Zonta M, Angulo MC, Gobbo S, Rosengarten B, Hossmann KA, Pozzan T, Carmignoto G: **Neuron-to-astrocyte signaling is central to the dynamic control of brain microcirculation.** *Nat Neurosci* 2003, **6**:43-50.
- Takano T, Tian GF, Peng W, Lou N, Libionka W, Han X, Nedergaard M: **Astrocyte-mediated control of cerebral blood flow.** *Nat Neurosci* 2006, **9**:260-267.
- Petzold GC, Albeanu DF, Sato TF, Murthy VN: **Coupling of neural activity to blood flow in olfactory glomeruli is mediated by astrocytic pathways.** *Neuron* 2008, **58**:897-910.
- Metea MR, Newman EA: **Glial cells dilate and constrict blood vessels: a mechanism of neurovascular coupling.** *J Neurosci* 2006, **26**:2862-2870.
- Filosa JA, Bonev AD, Nelson MT: **Calcium dynamics in cortical astrocytes and arterioles during neurovascular coupling.** *Circ Res* 2004, **95**:e73-81.
- Mulligan SJ, MacVicar BA: **Calcium transients in astrocyte endfeet cause cerebrovascular constrictions.** *Nature* 2004, **431**:195-199.
- Li A, Xi Q, Umstot ES, Bellner L, Schwartzman ML, Jaggar JH, Leffler CW: **Astrocyte-derived CO is a diffusible messenger that mediates glutamate-induced cerebral arteriolar dilation by activating smooth muscle cell KCa channels.** *Circ Res* 2008, **102**:234-241.
- Lou HC, Edvinsson L, MacKenzie ET: **The concept of coupling blood flow to brain function: revision required?** *Ann Neurol* 1987, **22**:289-297.
- Mintun MA, Vlassenko AG, Rundle MM, Raichle ME: **Increased lactate/pyruvate ratio augments blood flow in physiologically activated human brain.** *Proc Natl Acad Sci USA* 2004, **101**:659-664.
- Gordon GR, Choi HB, Rungta RL, Ellis-Davies GC, MacVicar BA: **Brain metabolism dictates the polarity of astrocyte control over arterioles.** *Nature* 2008, **456**:745-749.
- Anderson CM, Nedergaard M: **Astrocyte-mediated control of cerebral microcirculation.** *Trends Neurosci* 2003, **26**:340-344.
- Liu X, Li C, Falck JR, Roman RJ, Harder DR, Koehler RC: **Interaction of nitric oxide, 20-HETE, and EETs during functional hyperemia in whisker barrel cortex.** *Am J Physiol Heart Circ Physiol* 2008, **295**:H619-H631.
- Blanco VM, Stern JE, Filosa JA: **Tone-dependent vascular responses to astrocyte-derived signals.** *Am J Physiol Heart Circ Physiol* 2008, **294**:H2855-H2863.
- Fleming I: **Cytochrome P450 and vascular homeostasis.** *Circ Res* 2001, **89**:753-762.
- Iadecola C: **Regulation of the cerebral microcirculation during neural activity: is nitric oxide the missing link?** *Trends Neurosci* 1993, **16**:206-214.
- Hein TW, Xu W, Kuo L: **Dilation of retinal arterioles in response to lactate: role of nitric oxide, guanylyl cyclase, and ATP-sensitive potassium channels.** *Invest Ophthalmol Vis Sci* 2006, **47**:693-699.
- Kasischke KA, Vishwasrao HD, Fisher PJ, Zipfel WR, Webb WW: **Neural activity triggers neuronal oxidative metabolism followed by astrocytic glycolysis.** *Science* 2004, **305**:99-103.
- Pellerin L, Magistretti PJ: **Glutamate uptake into astrocytes stimulates aerobic glycolysis: a mechanism coupling neuronal activity to glucose utilization.** *Proc Natl Acad Sci USA* 1994, **91**:10625-10629.
- Swanson RA, Morton MM, Sagar SM, Sharp FR: **Sensory stimulation induces local cerebral glycogenolysis: demonstration by autoradiography.** *Neuroscience* 1992, **51**:451-461.
- Haglund MM, Meno JR, Hochman DW, Ngai AC, Winn HR: **Correlation of intrinsic optical signal, cerebral blood flow, and evoked potentials during activation of rat somatosensory cortex.** *J Neurosurg* 2008, **109**:654-663.
- Morii S, Ngai AC, Ko KR, Winn HR: **Role of adenosine in regulation of cerebral blood flow: effects of theophylline during normoxia and hypoxia.** *Am J Physiol* 1987, **253**:H165-H175.
- Ruth VJ, Park TS, Gonzales ER, Gidday JM: **Adenosine and cerebrovascular hyperemia during insulin-induced hypoglycemia in newborn piglet.** *Am J Physiol* 1993, **265**:H1762-H1768.
- Miekisiak G, Kulik T, Kusano Y, Kung D, Chen JF, Winn HR: **Cerebral blood flow response in adenosine 2a receptor knockout mice during transient hypoxic hypoxia.** *J Cereb Blood Flow Metab* 2008, **28**:1656-1664.
- Shi Y, Liu X, Gebremedhin D, Falck JR, Harder DR, Koehler RC: **Interaction of mechanisms involving epoxyeicosatrienoic acids, adenosine receptors, and metabotropic glutamate receptors in neurovascular coupling in rat whisker barrel cortex.** *J Cereb Blood Flow Metab* 2008, **28**:111-125.
- Anderson CM, Swanson RA: **Astrocyte glutamate transport: review of properties, regulation, and physiological functions.** *Glia* 2000, **32**:1-14.
- Shulman RG, Hyder F, Rothman DL: **Cerebral energetics and the glycogen shunt: neurochemical basis of functional imaging.** *Proc Natl Acad Sci USA* 2001, **98**:6417-6422.
- Rouach N, Koulakoff A, Abudara V, Willecke K, Giaume C: **Astroglial metabolic networks sustain hippocampal synaptic transmission.** *Science* 2008, **322**:1551-1555.